

NO-A102 908

FUNCTIONAL RELATIONSHIPS BETWEEN RISKY AND RISKLESS  
MULTIATTRIBUTE UTILIT.. (U) UNIVERSITY OF SOUTHERN  
CALIFORNIA LOS ANGELES SOCIAL SCIENCE R..


1/1


UNCLASSIFIED

D VON WINTERFELDT DEC 79 SSRI-RR-79-3

F/G 12/4

NL





(1)



JSC

# social science research institute

## RESEARCH REPORT

### FUNCTIONAL RELATIONSHIPS BETWEEN RISKY AND RISKLESS MULTIATTRIBUTE UTILITY FUNCTIONS

Detlof von Winterfeldt

#### Sponsored By:

Defense Advanced Research Projects Agency  
Department of the Navy  
Office of Naval Research

#### Monitored By:

Department of the Navy  
Office of Naval Research  
Pasadena Branch Office  
Contract No. N00014-79-C-0038

Approved for Public Release;  
Distribution Unlimited;  
Reproduction in Whole or in Part is  
Permitted for Any Use of the U.S. Government

DECEMBER, 1979

SSRI RESEARCH REPORT 79-3

DTIC  
ELECTE  
JUL 29 1987  
S D  
CD

AD-F  
UNIVERSITY OF SOUTHERN CALIFORNIA

DISTRIBUTION STATEMENT

Approved for public release;  
Distribution Unlimited

TECHNICAL REPORT SSRI 79-3

FUNCTIONAL RELATIONSHIPS BETWEEN RISKY AND  
RISKLESS MULTIATTRIBUTE UTILITY FUNCTIONS

by

Detlof von Winterfeldt

Sponsored by

Defense Advanced Research Projects Agency

December, 1979

SOCIAL SCIENCE RESEARCH INSTITUTE  
University of Southern California  
Los Angeles, California 90007

|                    |  |
|--------------------|--|
| Accession For      |  |
| NTIS CRA&I         | <input checked="checked" type="checkbox"/> |
| DTIC TAB           | <input type="checkbox"/>                   |
| Unannounced        | <input type="checkbox"/>                   |
| Justification      |  |
| By                 |  |
| Distribution/      |  |
| Availability Codes |  |
| Dist               | Avail and/or Special                       |
| A-1                |  |



## ABSTRACT

Expected utility theory and conjoint measurement theory form two major classes of models and assessment procedures to construct multi-attribute utility functions. In conjoint measurement theory a value function  $v$  is constructed which preserves preferences among riskless multi-attributed outcomes. The risky utility function  $u$ , constructed in the framework of expected utility theory, also preserves such riskless preferences. In addition,  $u$  is an appropriate guide for decisions under uncertainty since its expectation preserves risky preferences among gambles. Since both  $u$  and  $v$  are order preserving functions, they must be related by a strictly increasing transformation. However,  $u$  and  $v$  need not coincide or be related through any special functional forms, unless some simple decomposition forms are assumed. More restricted functional relationships obtain, if  $u$  and  $v$  are assumed to be either additive or multiplicative. In particular,  $u$  can be shown to be linearly, logarithmically, or exponentially related to  $v$ , depending on which function is additive and which is multiplicative. The paper proves such functional relationships based on the theory of functional equations, and techniques are described to assess the parameters of these functions. The results are discussed from a behavioral standpoint of interpreting the form and shape of multi-attribute utility functions and from a practical standpoint of simplifying multi-attribute utility assessment.

## TABLE OF CONTENTS

|  | <u>Page</u> |
|--|-------------|
| ABSTRACT . . . . .                           | ii          |
| LIST OF FIGURES. . . . .                     | iv          |
| ACKNOWLEDGEMENT. . . . .                     | v           |
| DISCLAIMER . . . . .                         | vi          |
| INTRODUCTION . . . . .                       | 1           |
| DEFINITIONS AND PRELIMINARY RESULTS. . . . . | 6           |
| BEHAVIORAL AND APPLIED IMPLICATIONS. . . . . | 30          |
| REFERENCES . . . . .                         | 60          |

## LIST OF FIGURES

|    |  | <u>Page</u> |
|----|--|-------------|
| 1. | Global Properties of Functions Relating $u$ and $v$ if $u$ is Multiplicative and $v$ is Additive (Standardization conventions as in text)..... | 25          |
| 2. | Local Properties of Function Relating $u$ and $v$ if $u$ is Multiplicative and $v$ is Additive (Standardization conventions as in text).....   | 26          |
| 3. | Global Properties of Functions Relating $u$ and $v$ if Both are Multiplicative (Standardization conventions for $k > 0$ as in text).....       | 28          |
| 4. | Local Properties of Functions Relating $u$ and $v$ if Both are Multiplicative (Standardization conventions for $k > 0$ as in text).....        | 29          |
| 5. | Global Properties of Functions Relating $u$ and $v$ if Both are Multiplicative (Standardization conventions for $k < 0$ as in text).....       | 31          |
| 6. | Local Properties of Functions Relating $u$ and $v$ if Both are Multiplicative (Standardization conventions for $k > 0$ as in text).....        | 32          |

# ACKNOWLEDGEMENT

This research was supported by the Advanced Research Projects Agency of the Department of Defense, and was monitored by the Office of Naval Research under Contract No. N00014-79-C-0038.

# DISCLAIMER

The views and conclusions contained in this document are those of the author and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the Advanced Research Projects Agency or of the United States Government.

## INTRODUCTION

Multiattribute preferences can be modeled in two fundamentally different ways, depending on whether the decision problem involves uncertainties or not. The first modeling approach is based on conjoint measurement theory (see Krantz, Luce, Suppes, and Tversky, 1971) which applies to the problem of modeling riskless preferences for multiattributed outcomes. Conjoint measurement theory specifies the conditions under which a riskless value function  $v$  can be constructed which preserves the preference order among multiattributed outcomes and which can be expressed as a simple aggregate of single attribute value functions  $v_i$ . The best known conjoint measurement model forms are the additive and the multiplicative models:

$$v(x) = \sum_{i=1}^n v_i(x_i) \quad (1)$$

and

$$v(x) = \prod_{i=1}^n v_i(x_i), \quad (2)$$

where  $x \in X$  is a multiattribute outcome,  $x_i$  is the level of  $x$  in the  $i$ -th attribute  $X_i$ ,  $v_i$  is the  $i$ -th single attribute value function, and  $v$  is the overall value function. Other simple polynomial forms of  $v$  have been developed in Krantz and Tversky (1970) and Krantz et al (1971). Although  $v$  is an appropriate guide for preferences among sure things, nothing in conjoint measurement theory guarantees that the expectation of  $v$  is appropriate for selecting among gambles with multiattributed outcomes.

The second modeling approach, expected utility theory, explicitly addresses the problem of decision making under risk. Based on v. Neumann and Morgenstern's (1947) work which was later extended by Savage (1954) and others, this theory provides the rationale for constructing a utility function  $u$  which preserves the preferences for riskless outcomes and at the same time its expectation preserves the preferences among gambles for such outcomes. Applied to the multi-attribute situation, several decomposition forms of  $u$  have been developed (see Keeney and Raiffa, 1976). The additive and the multiplicative forms are the best known:

$$u(x) = \sum_{i=1}^n u_i(x_i), \quad (3)$$

$$1 + k u(x) = \prod_{i=1}^n [1 + k u_i(x_i)]. \quad (4)$$

It is relatively trivial to establish the fact that  $u$  and  $v$  are related by a strictly increasing transformation (see Raiffa, 1969; Krantz et al, 1971; Keeney and Raiffa, 1976). But neither conjoint measurement theory, nor expected utility theory by themselves provide a rationale for any specific functional relationships between  $u$  and  $v$ . In principle, the shape and aggregation rule of  $u$  and  $v$  can be quite different. For example,  $v$  may be additive, while  $u$  is multiplicative or non-decomposable at all. All  $v_i$ 's may be linear, while all  $u_i$ 's may be non-linear, etc.

Establishing closed form functional relationships between  $u$  and  $v$  is, however, possible when special decomposition forms such as (1) - (4) are assumed. Scattered throughout the multiattribute literature

are results which relate utility and value functions by some specific class of transformation, e.g., exponential or logarithmic functions (see Pollak, 1967; Krantz et al, 1971; Keeney and Raiffa, 1976; Dyer and Sarin, 1977). Keeney and Raiffa, for example, give a proof outline to show that an additive value function  $v$  and a multiplicative utility function  $u$  must be related exponentially:

$$u(x) = b \exp\{a v(x)\}. \quad (5)$$

Another way to relate  $u$  and  $v$  is by the uniqueness theorems of their respective measurement theoretic representations. For example, an additive conjoint measurement function  $v$  is unique up to a positive linear transformation. Consequently, any other additive and order preserving function  $v'$  must be related to  $v$  by

$$v'(x) = a \cdot v(x) + b. \quad (6)$$

for some  $a > 0, b$ . In particular, an additive utility function  $u$  should be related to an additive value function by a positive linear transformation.

Functional forms such as (5) and (6) have been proven by a variety of mathematical methods (differentiation methods in Pollak, constructive algebraic proofs in Krantz et al, risk attitude arguments in Keeney and Raiffa, and uniqueness theorems in Dyer and Sarin). No common framework or integrated presentation of these functional relations has been developed yet. This paper attempts to provide such an integrative framework through the theory of functional equations.

The interest in parametric functional relationships such as (5) and (6) is not merely theoretical. For the practitioner who has to assess multiattribute utility functions in an applied context such functional forms provide tools for simplifying and cross-checking utility and value functions. Simplifications in the construction of  $u$  are possible, by first constructing  $v$  and then exploring the functional form relating  $u$  and  $v$  through its parametric properties. Such a two step construction process has several advantages. First,  $v$  can usually be approximated by simple techniques such as Edwards' SMART procedure (1977), which involves similar judgmental processes as the theoretically feasible construction process of dual standard sequences (see Krantz et al, 1971). Second, the two step procedure avoids lengthy and tedious lottery assessments in the construction of  $u$  by assessing the parameters of the function  $u = h(v)$  through a few simple questions involving risky outcomes.

Constructing the utility function  $u$  in the two step process has also theoretical advantages. The value function  $v$  and its single attribute functions  $v_i$  express purely riskless preference characteristics, such as marginally decreasing value, complementarity or substitution phenomena between attributes. Such considerations are usually compounded in  $u$  and its single attribute functions  $u_i$  with the risk attitudes of the decision maker. For example, it is quite conceivable that a single attribute utility function  $u_i$  appears risk neutral (linear) because marginal decreasing value is compensated by risk proneness. The two step construction process uncovers such anomalies by identifying riskless value aspects in the value function, and by incorporating

"pure" risk considerations in the transformation  $u = h(v)$ .

The next section will show that  $u$  and  $v$  must be related by a strictly increasing transformation  $h$ , which under further technical assumptions must be continuous. Then four possible functional equations will be investigated and solved.

$$u = \sum_{i=1}^n u_i = h \left[ \sum_{i=1}^n v_i \right] = h(v), \quad (7)$$

$$1+ku = \prod_{i=1}^n (1+ku_i) = h \left[ \sum_{i=1}^n v_i \right] = h(v), \quad (8)$$

$$u = \sum_{i=1}^n u_i = h \left[ \prod_{i=1}^n v_i \right] = h(v), \quad (9)$$

$$1+ku = \prod_{i=1}^n (1+ku_i) = h \left[ \prod_{i=1}^n v_i \right] = h(v). \quad (10)$$

The solutions consist of a reduction to one of the fundamental Cauchy type functional equations (see Aczél, 1966). For certain standardizations of  $u$  and  $v$  these solutions turn out to be very simple, namely  $u = v$  for (7),  $1+ku = (1+k)^v$  for (8),  $u = \ln v$  for (9) and  $1 + ku = (\text{sgnk})(\text{sgnv})|v|^{\ln[1+(\text{sgnk}) \cdot k]}$  for (10). Subsequent to proving these solutions, some behavioral and practical implications will be discussed.\*

\* Readers who are willing to accept the above solutions and do not want to bother with the rather technical proofs, are recommended to skip the following two sections.

### Definitions and Preliminary Results

The value and utility functions  $v$  and  $u$  will be defined on the product set  $X = X_1 \times X_2 \times \dots \times X_i \times \dots \times X_n$ .  $x$  and  $y$  are typical elements of  $X$ .  $x_i$  and  $y_i$  characterize their respective levels in attribute  $X_i$ .  $P$  is the set of simple probability distributions over  $X$  with typical elements  $p, q \in P$ . A transitive and connected order relation  $\succsim$  is defined on  $P$  with  $p \succsim q$  being interpreted as "p is preferred to or indifferent to q." By considering degenerate probability distributions  $\succsim$  can be reduced to  $X$ . Thus  $p \succsim q$  where  $p(x) = q(y) = 1$  for some  $x, y \in X$  is written as  $x \succsim y$ , and interpreted as "x is preferred to or indifferent to y."  $u$  and  $v$  are defined as follows (for more details, see Fishburn, 1970):

Definition 1 A function  $u: X \rightarrow \mathbb{R}$  is a (v. Neumann and Morgenstern) utility function if for all  $p, q \in P$

$$\begin{aligned} P &\succsim q \\ \text{if and only if} & \\ E(u | p) &\geq E(u | q) \end{aligned} \tag{11}$$

where  $E(u | \cdot)$  denotes the expectation of  $u$  with respect to some probability distribution. If  $u$  can be expressed as in (3) it is called additive. If  $u$  can be expressed as in (4) it is called multiplicative.

Note: From Definition 1 and the reduction of  $\succsim$  to  $X$  it follows that  $u$  is order preserving, i.e., for all  $x, y \in X$

$$\begin{aligned} x &\succsim y \\ \text{if and only if} & \\ u(x) &\geq u(y) \end{aligned} \tag{12}$$

Definition 2 A function  $v: X \rightarrow \text{Re}$  is a (conjoint measurement) value function, if for all  $x, y \in X$

$$\begin{aligned} x \succsim y \\ \text{if and only if} \\ v(x) \geq v(y). \end{aligned} \tag{13}$$

If  $v$  can be expressed as in (1) it is called additive.

If  $v$  can be expressed as in (2), it is called multiplicative.

The axioms and representation theorems leading to definitions 1 and 2 are of no special interest here. Axiomatic foundations of utility functions can be found in v. Neumann and Morgenstern (1947), Savage (1954), and Fishburn (1970). Axioms for the additive and multiplicative decomposition forms of  $u$  are presented in Keeney and Raiffa (1976) and Fishburn (1970). Axiomatic foundations of conjoint measurement value functions are given in Krantz et al (1971), including the additive and the multiplicative forms as well as other simple polynomials.

Besides the assumptions implicit in definitions 1 and 2, the solutions to the functional equations (7)-(10) require that  $h$  be continuous. Since continuity is a rather abstract concept, the more natural assumption will be made in the following that  $u$  and  $v$  are defined onto some convex subset of  $\text{Re}$ , labelled  $I_u$  and  $I_v$  respectively. From the definitions of  $u$  and  $v$  and from this onto property it follows that  $h$  must be strictly increasing and continuous. Lemma 1 formalizes this implication.

Lemma 1 Assume that  $u: X \xrightarrow{\text{onto}} I_u$  is a utility function, where  $I_u$  is a convex subset of  $\text{Re}$ . Assume further that  $v: X \xrightarrow{\text{onto}} I_v$  is a value function where  $I_v$  is a convex subset of  $\text{Re}$ . Then there exists a function  $h: I_v \xrightarrow{\text{onto}} I_u$  which is strictly increasing and continuous.

Proof. The proof is trivial except for continuity. Define  $h$

by  $u(x) = h[v(x)]$  for all  $x \in X$ . From the equality part of (12) and (13) it follows that  $h$  is well defined. From the inequality part of (12) and (13) it follows that  $h$  is strictly increasing.  $h$  is onto since both  $u$  and  $v$  are onto.

To prove continuity, consider the contrary. Then there exists at least one point  $r \in I_v$  at which  $h$  is discontinuous. By the onto property of  $v$  and the definition of  $h$ ,  $h(r)$  is defined. Consider first the case where  $r$  is not a boundary point of  $I_v$ , and define right and left hand limits as

$$\begin{aligned} \lim_{t \rightarrow r^-} h(t) &= L, \\ \lim_{t \rightarrow r^+} h(t) &= R. \end{aligned} \tag{14}$$

By the assumption that  $r$  is not a boundary point, we can define a sufficiently small  $\epsilon$  such that  $h(r+\epsilon)$  and  $h(r-\epsilon)$  exist. By the fact that  $h$  is strictly increasing

$$h(r-\epsilon) < L \leq h(r) \leq R < h(r+\epsilon). \tag{15}$$

This establishes boundaries for  $L$  and  $R$  and shows that both limits must exist. By the assumed discontinuity at  $r$ , however, at least one of the weak inequalities in (15) must be strict. By the convexity of  $I_u$  we therefore can find an  $a$  such that  $m = aR + (1-a)L \neq h(r)$ ;  $m \in I_u$ . However, by (15) there exists no  $s \in I_v$  such that  $h(s) = m$ , thus contradicting the onto property of  $h$ . Consequently,  $h$  must be continuous, at  $r$ .

A similar argument can be made for the cases where  $r$  is either a lower or an upper boundary point of  $I_v$  by only considering right hand or left hand limits. In each of these cases the assumption that  $h$  is discontinuous leads to a violation of the onto property of  $h$ . Thus  $h$  must be continuous everywhere in  $I_v$ .

In solving the functional equations (7)-(10) it is often convenient to transform  $u$  and  $v$ . The following simple lemmas state which transformations are admissible in the sense that they do not "destroy" any property of a value on utility functions. These lemmas are actually the necessary parts of the uniqueness theorems in expected utility theory and conjoint measurement theory. Since the proofs are very simple, only proof notes are given for each lemma.

Lemma 2 If  $v$  is an additive value function with additive terms  $v_i$ ,

then  $v' = av + b$ ,  $a > 0$ , is also an additive value function with additive terms  $v_i' = av_i + b_i$ ,  $\sum_{i=1}^n b_i = b$ .

Proof note. Since  $a$  is positive,  $v'$  is order preserving. Also,  $v'$  can be written as an additive function by using  $v_i'$ . Therefore  $v'$  is an additive value function.

Lemma 3 If  $v$  is a multiplicative value function with multiplicative

terms  $v_i$ , then  $v' = (\text{sgn } v) \cdot b \cdot |v|^a$ ,  $a, b > 0$ , is also a multiplicative value function with multiplicative terms  $v_i' = (\text{sgn } v_i) \cdot b_i |v_i|^a$ , where  $\prod_{i=1}^n b_i = b$ .

Proof note. Since  $a$  and  $b$  are positive,  $v'$  is order preserving.

Also,  $v'$  can be written as a multiplicative function using terms  $v_i'$ .

The sign conditions make sure that  $v$  and  $v'$  have identical signs everywhere.

Lemma 4 If  $u$  is an additive utility function with additive terms  $u_i$ , then  $u' = au + b$  is also an additive utility function with additive terms  $u_i' = au_i + b_i$ , where  $\sum_{i=1}^n b_i = b$ .

Proof note. Since  $a$  is positive,  $u'$  is order preserving over  $X$ . Since  $E(u' | \cdot) = a E(u | \cdot) + b$ , the expectation of  $u'$  is also order preserving over  $P$ . Finally  $u'$  can be expressed as an additive function using the  $u_i'$ 's.

Lemma 5 If  $k > 0$  and  $1+ku$  is a multiplicative utility function with multiplicative terms  $1+ku_i$ , then  $u' = a(1+ku)$ ,  $a > 0$  is also a multiplicative utility function with multiplicative terms  $u_i' = a_i(1+ku_i)$ ,  $\prod_{i=1}^n a_i = a$ . If  $k < 0$  and  $-(1+ku)$  is a multiplicative utility function with multiplicative terms  $(1+ku_i)$ , then  $u' = -a(1+ku)$ ,  $a > 0$ , is also a multiplicative utility function with multiplicative terms  $a_i(1+ku_i)$ ,  $\prod_{i=1}^n a_i = -a$ .

Proof note. The two cases  $k > 0$  and  $k < 0$  must be considered separately to choose appropriate signs of the constants. (The case  $k = 0$  is equivalent to  $u$  being additive; thus Lemma 3 applies.) If  $k > 0$ , and  $u$  is the original utility function, then  $1+ku$  and  $u'$  are also utility functions since  $a$  is positive and since the expectation operator is linear. Also,  $u'$  can be expressed multiplicatively by using the  $u_i'$ 's. The sign of the  $a_i$ 's is arbitrary as long as their product is positive, thus maintaining the order preserving property of  $u'$ . If  $k < 0$ , and  $u$  is the original utility function,  $1+ku$  produces an inverse ordering. Thus  $-(1+ku)$  is again a utility function of the form  $-(1+ku) = -\prod_{i=1}^n (1+ku_i)$ . Since  $a$  is positive  $u'$  also is a utility function, which can be expressed multiplicatively by using  $u_i'$ . The scaling of the  $a_i$ 's

guarantees that the product of the single attribute  $u_i$ 's is again negative.

A point worth noting here is that the multiplicative value functions could be transformed exponentially while still maintaining order preserving properties and multiplicativeness. Such transformations are not admissible for utility functions since they would destroy their expectation property.

### The Basic Functional Equations and Their Solutions

With these results as a background, functional equations (7)-(10) will now be solved by a reduction to fundamental Cauchy type equations with some simple solution. In the simplest case, both  $u$  and  $v$  are additive:

Theorem 1 Assume that  $u$  and  $v$  are additive as in (1) and (3). Then there exist real numbers  $a > 0$ ,  $b$  such that

$$u = av + b \quad (14)$$

Proof. The uniqueness of  $u$  and  $v$  can be used to prove the theorem (see Dyer and Sarin, 1977). However, uniqueness is itself a rather difficult property to establish, and its proof is usually hidden in the constructive algebraic proofs of the representation theorems leading up to the definition of utility and value functions. This proof therefore uses the more fundamental route of reducing (7) to Cauchy's equation  $h(x+y) = h(x) + h(y)$ .

Consider some arbitrary  $x^0 = (x_1^0, x_2^0, \dots, x_i^0, \dots, x_n^0)$  with  $v_i(x_i^0) = v_i^0$ ,  $v(x^0) = v^0$ ,  $u_i(x_i^0) = u_i^0$ , and  $u(x^0) = u^0$ . Define transformed value and utility functions by

$$v_i' = v_i - v_i^0, \quad (15)$$

$$v' = \sum_{i=1}^n v_i', \quad (16)$$

$$u_i = u_i - u_i^0, \quad (17)$$

$$u' = \sum_{i=1}^n u_i'. \quad (18)$$

By Lemma 2  $v'$  is an additive value function, and by Lemma 4  $u'$  is an additive utility function. Furthermore if  $v$  and  $u$  are defined onto some real interval, so are  $v'$  and  $u'$ . Therefore, Lemma 1 applies and there exists a strictly increasing continuous function  $h$  with

$$\sum_{i=1}^n u_i'(x_i) = h\left[\sum_{i=1}^n v_i'(x_i)\right]. \quad (19)$$

Let  $x_j = x_j^0$  for all  $j \neq i$ . By (15) and (17)  $v_j'(x_j^0) = 0$  and  $u_j'(x_j^0) = 0$ , thus (19) simplifies to

$$u_i'(x_i) = h[v_i'(x_i)]. \quad (20)$$

Substituting the right hand side of (20) into (19) gives

$$\sum_{i=1}^n h[v_i'(x_i)] = h\left[\sum_{i=1}^n v_i'(x_i)\right]. \quad (21)$$

In particular, let  $x_i = x_i^0$  for  $i = 3, 4, \dots, n$ . This reduces (21) to

$$h[v_1'(x_1)] + h[v_2'(x_2)] = h[v_1'(x_1) + v_2'(x_2)]. \quad (22)$$

(22) is Cauchy's fundamental functional equation  $h(x) + h(y) = h(x + y)$  (see Aczél, 1966), which for continuous  $h$  has the non-trivial solution

$$h(r) = ar. \quad (23)$$

Substituting (23) into (20) and (19) gives

$$u_i'(x_i) = av_i'(x_i), \quad (24)$$

$$\sum_{i=1}^n u_i'(x_i) = a \sum_{i=1}^n v_i'(x_i), \quad (25)$$

and  $u' = a v'. \quad (25)$

Resubstituting  $u$  and  $u$  for  $u'$  and  $v'$  gives the desired result

$$u = av + b,$$

where  $b = u^0 - av^0$ .  $a > 0$  follows directly from the fact that  $h$  is strictly increasing.

**Theorem 2** Assume that  $(1+ku)$  is multiplicative as in (4) and that  $v$  is additive as in (1). Then there exist  $a > 0$ ,  $b > 0$  such that

$$1+ku = b \exp \{(\text{sgn } k) a v\} \quad (26)$$

**Proof.** The proof is based on a reduction of (8) to the functional equation  $h(xy) = h(x) + h(y)$ . Again, consider some arbitrary

$x^0 = (x_1^0, x_2^0, \dots, x_i^0, \dots, x_n^0)$  and define  $v_i^0$ ,  $v^0$ ,  $u_i^0$  and  $u^0$ ,  $v_i'$  and  $v'$  as in the proof of Theorem 1. Define the transformed utility

function  $u'$  by

$$u_i' = \frac{1}{(1+ku_i^0)} (1+ku_i), \quad (27)$$

$$u' = \begin{cases} \prod_{i=1}^n u_i' & \text{if } k > 0, \\ -\prod_{i=1}^n u_i' & \text{if } k < 0. \end{cases} \quad (28)$$

$v'$  is an additive value function (Lemma 2). To establish that  $u'$  is a multiplicative utility function, observe that  $[(1+ku_i(x_i))] > 0$  for all  $i = 1, 2, \dots, n$  and  $x_i \in X_i$ . For if there was a  $y_i$  such that  $1+ku_i(y_i) \leq 0$ , then reversals of preferences must occur or all elements in  $X$  must be indifferent for that value of  $y_i$ . (See also Fishburn and Keeney, 1974.) Neither case is compatible with the additive value function. Therefore the sign scaling conditions of Lemma 5 are valid, and  $u'$  is a multiplicative utility function. In addition, since  $v$  and  $u$  are defined onto some convex subset of  $\mathbb{R}^n$ , so are  $u'$  and  $v'$ . Therefore Lemma 1 applies, and there exists a strictly increasing continuous function  $h$  such that

$$\prod_{i=1}^n u_i'(x_i) = h[(\text{sgn } k) \sum_{i=1}^n v_i'(x_i)]. \quad (29)$$

(Note: (29) is the compact form of the two functional equations  $u' = h(v')$  for  $k > 0$  and  $-u' = h(-v')$  for  $k < 0$ ). Evaluating (29) at  $x_j = x_j^0$  for  $j \neq i$  and noting that  $u_j'(x_j^0) = 1$  and  $v_j'(x_j^0) = 0$  reduces (29) to

$$u_i'(x_i) = h[(\text{sgn } k) v_i'(x_i)]. \quad (30)$$

Substituting the right hand side of (30) into (29) gives

$$\prod_{i=1}^n [h(\text{sgn } k) v_i'(x_i)] = h[(\text{sgn } k) \sum_{i=1}^n v_i'(x_i)]. \quad (31)$$

In particular, let  $x_i = x_i^0$  for  $i = 3, 4, \dots, n$ . This reduces (31) to

$$\begin{aligned} h[(\text{sgn } k) v_1(x_1)] h[(\text{sgn } k) v_2(x_2)] = \\ h\{[(\text{sgn } k) v_1(x_1)] + [(\text{sgn } k) v_2(x_2)]\}. \end{aligned} \quad (32)$$

(32) is the fundamental functional evaluation  $h(xy) = h(x + y)$  (see Aczel, 1961), which for continuous  $h$  has the non-trivial solution

$$h(r) = \exp(a r). \quad (33)$$

Since  $h$  is strictly increasing,  $a > 0$ .

Substituting (33) into the original functional equations (29) and (30) gives

$$u_i'(x_i) = \exp[(\text{sgn } k) a v_i'(x_i)], \quad (34)$$

$$\text{and} \quad \prod_{i=1}^n u_i'(x_i) = \exp[(\text{sgn } k) a \sum_{i=1}^n v_i'(x_i)]. \quad (35)$$

Resubstituting  $u_i$  and  $v_i$  for  $u_i'$  and  $v_i'$  gives the desired results:

$$1+ku_i(x_i) = (1+ku_i^0) \exp \{(\operatorname{sgn} k) a [v_i(x_i) - v_i^0]\}, \quad (36)$$

$$\text{and } \prod_{i=1}^n [1+ku_i(x_i)] = \prod_{i=1}^n (1+ku_i^0) \cdot \exp \{(\operatorname{sgn} k) a \sum_{i=1}^n [v_i(x_i) - v_i^0]\}, \quad (37)$$

which, after appropriate definitions of constants becomes

$$1 + ku = b \exp [(\operatorname{sgn} k) a v], \quad b, a > 0. \quad (38)$$

Another way of writing (38) is somewhat more instructive:

$$u = (\operatorname{sgn} k) b' \exp \{(\operatorname{sgn} k) a v\} + c. \quad (39)$$

This form shows that  $k$  directly controls the shape of the exponential transformation. In utility theoretic terms, if  $k < 0$  then  $h$  is a constantly risk averse transformation, if  $k > 0$  it is a constantly risk seeking transformation. This result was previously shown by Keeney and Raiffa, 1976, (p.331).

The next theorem reverses the roles of  $u$  and  $v$ :

**Theorem 3** If  $u$  is additive as in (3) and  $v$  is multiplicative as in (2), then there exist  $a > 0$ ,  $b$ , such that

$$u = a \ln v + b, \quad (40)$$

where  $v > 0$ .

**Proof.** The proof reduces (9) to the functional equation  $h(x) + h(y) = h(xy)$ . Consider some arbitrary  $x^0 = (x_1^0, x_2^0, \dots, x_i^0, \dots, x_n^0)$  and define  $v_i^0, v^0, u_i^0, u^0, u_i'$  and  $u'$  as in Theorem 1.

The transformed value function is defined by

$$v_i' = \frac{v_i}{v^0} \quad (41)$$

$$v' = \frac{v}{v^0} \quad (42)$$

$u'$  is an additive utility function (Lemma 4). To establish that  $v_i'$  and  $v'$  exist and represent a multiplicative value function, it suffices to show that for all  $i$ ,  $x_i \in X_i$ ,  $v_i(x_i) > 0$ . If there was an  $y_i$  such that  $v_i(y_i) \leq 0$ , then reversals of preferences must occur among elements in  $X$  or all elements in  $X$  must be indifferent for that value of  $y_i$ . (See also Fishburn and Keeney, 1974.) Both cases are incompatible with the additive utility function  $u$ . Therefore (41) and (42) are positive transformations and  $v'$  is a multiplicative value function (Lemma 3). Since both  $u$  and  $v$  are defined onto some convex subset of  $\mathbb{R}^n$ , so are  $u'$  and  $v'$ . Therefore there exist a strictly increasing continuous function  $h$  such that

$$\sum_{i=1}^n u_i'(x_i) = h \left[ \prod_{i=1}^n v_i'(x_i) \right]. \quad (43)$$

Evaluating (43) at  $x_j = x_j^0$  for  $j \neq i$  and, noting that  $u_j'(x_j^0) = 0$  and  $v_j(x_j^0) = 1$ , (43) can be reduced to

$$u_i'(x_i) = h[v_i'(x_i)]. \quad (44)$$

Substituting (44) into (43) gives

$$\sum_{i=1}^n h[v_i'(x_i)] = h\left[\prod_{i=1}^n v_i'(x_i)\right]. \quad (45)$$

In particular, letting  $x_i = x_i^0$  for  $i = 3, 4, \dots, n$  reduces (45) to

$$h[v_1'(x_1)] + h[v_2'(x_2)] = h[v_1'(x_1)v_2'(x_2)]. \quad (46)$$

(46) is the functional equation  $h(x) + h(y) = h(xy)$  (see Aczél, 1966) which for continuous  $h$  and positive  $r$  has the solution

$$h(r) = a \ln r, \quad (47)$$

where  $a > 0$ , since  $h$  is strictly increasing.

Substituting (47) into the original functional equations (44) and (45) gives

$$u_i'(x_i) = a \ln v_i'(x_i), \quad (48)$$

and

$$\sum_{i=1}^n u_i'(x_i) = a \ln \left[ \prod_{i=1}^n v_i'(x_i) \right]. \quad (49)$$

Resubstituting  $u_i$  and  $v_i$  for  $u_i'$  and  $v_i'$  gives the desired results:

$$u_i = a \ln \frac{v_i}{v_i^0} + u_i^0 \quad (50)$$

and

$$u = a \ln \frac{v}{v^0} + u^0, \quad (51)$$

which, after appropriate definitions of constants is

$$u = a \ln v + b. \quad (52)$$

Theorem 4 If both  $u$  and  $v$  are multiplicative as in (2) and (4), then there exist  $a, b > 0$  such that

$$1 + ku = (\text{sgn } k)(\text{sgn } v) b|v|^a \quad (53)$$

Proof. The proof consists of reducing functional equation (10) to the fundamental equation  $h(xy) = h(x)h(y)$ . Although the route will be similar to the previous theorems, the situation here is more general since sign reversals and null region are allowed. Again some  $x^0 = (x_1^0, x_2^0, \dots, x_i^0, \dots, x_n^0)$  is picked, but this time zero multipliers must be avoided, i.e.,  $1 + ku_i(x_i^0) \neq 0$ ;  $v_i(x_i^0) \neq 0$ . There is no problem with excluding zero multipliers, as long as there exist at least one  $x$  and one  $y$  in  $X$  such that  $x \succ y$ . Because if in one attribute the only realizable value  $z_i^0$  was a zero multiplier, all elements in  $X$  would be indifferent.

Let  $v_i^0, v^0, u_i^0, u^0$  be defined as in the previous theorem.

Transformed value and utility functions are defined as follows:

$$v_i' = \begin{cases} \frac{v_i}{v_i^0} & \text{if } v_i^0 > 0, \\ -\frac{v_i}{v_i^0} & \text{if } v_i^0 < 0. \end{cases} \quad (54)$$

$$v' = \prod_{i=1}^n v_i' \quad (55)$$

$$u_i' = \begin{cases} \frac{1+ku_i}{1+ku_i^0} & \text{if } 1+ku_i^0 > 0, \\ -\frac{1+ku_i}{1+ku_i^0} & \text{if } 1+ku_i^0 < 0. \end{cases} \quad (56)$$

$$u' = \begin{cases} \prod_{i=1}^n u_i' & \text{if } k > 0, \\ -\prod_{i=1}^n u_i' & \text{if } k < 0. \end{cases} \quad (57)$$

The above definitions guarantee that all single attribute transformations are positive, thus the sign scaling conditions of Lemmas 3 and 5 are met. Consequently  $u'$  and  $v'$  are again proper multiplicative value and utility functions. In addition, since  $u$  and  $v$  are defined onto some convex subset of  $\mathbb{R}^n$ , so are  $u'$  and  $v'$ . Thus Lemma 1 applies and there exists a strictly increasing continuous function  $h$  such that

$$\prod_{i=1}^n u_i'(x_i) = h[(\text{sgn } k) \prod_{i=1}^n v_i'(x_i)]. \quad (58)$$

(Note: (58) is the compact form of the two functional equations  $u' = h(v')$  for  $k > 0$  and  $-u' = h(-v')$  for  $k < 0$ .) Evaluating (58) at  $x_j = x_j^0$  for  $j \neq i$ , and noting that  $u_j'(x_j^0) = v_j'(x_j^0) = 1$  reduces (58) to

$$u_i'(x_i) = h[(\text{sgn } k) v_i'(x_i)]. \quad (59)$$

Substituting the right hand side of (59) into (58) yields

$$\prod_{i=1}^n h[(\operatorname{sgn} k) v_i'(x_i)] = h[(\operatorname{sgn} k) \prod_{i=1}^n v_i'(x_i)]. \quad (60)$$

In particular, let  $x_i = x_i^0$  for  $i=3,4,\dots, n$ . This reduces (60) to

$$\begin{aligned} h[(\operatorname{sgn} k) v_1(x_1)] h[(\operatorname{sgn} k) v_2(x_2)] = \\ h\{[(\operatorname{sgn} k) v_1(x_1)][(\operatorname{sgn} k) v_2(x_2)]\}. \end{aligned} \quad (61)$$

(62) is the fundamental functional equation  $h(x)h(y) = h(xy)$  (see Aczél, 1966) which, for continuous  $h$  has the nontrivial solution

$$h(r) = (\operatorname{sgn} r) |r|^a. \quad (62)$$

Since  $h$  must be strictly increasing  $a > 0$ . (See Aczél, 1966.)

Substituting (62) into the original functional equations (58) and (59) gives:

$$u_i'(x_i) = (\operatorname{sgn} k)[\operatorname{sgn} v_i'(x_i)]|v_i'(x_i)|^a \quad (63)$$

and 
$$\prod_{i=1}^n u_i'(x_i) = (\operatorname{sgn} k)[\operatorname{sgn} \prod_{i=1}^n v_i'(x_i)]|\prod_{i=1}^n v_i'(x_i)|^a. \quad (64)$$

Resubstituting  $u_i$  and  $v_i$  for  $u_i'$  and  $v_i'$  gives the desired results:

$$1+ku_i = (1+ku_i^0)(\operatorname{sgn} k)(\operatorname{sgn} \frac{v_i}{v_i^0})|v_i^0|^{-a} |v_i|^a, \quad (65)$$

$$\prod_{i=1}^n 1+ku_i = \prod_{i=1}^n (1+ku_i^0) (\text{sgn } k) (\text{sgn } \prod_{i=1}^n \frac{v_i}{v_i^0}) \left| \prod_{i=1}^n v_i^0 \right|^{-a} \left| \prod_{i=1}^n v_i \right|^a, \quad (66)$$

which, after appropriate definitions of signs and constant terms gives

$$1+ku = (\text{sgn } k)(\text{sgn } v) b |v|^a \quad (67)$$

where  $b, a > 0$ .

### Scaling Procedures and Examples

The solutions of the functional equations presented in Theorems 1-4 are unique up to the specification of two parameters. By Lemmas 2-5, however, we can transform value and utility functions using two free parameters. Thus it is possible to use standardization conventions in the construction of  $u$  and  $v$  such that both functions go through two arbitrary fixed points. Such conventions will "consume" the two free parameters in the functional equations and produce special forms of functional equations which do not depend any more on  $a$  and  $b$ .

For the case in which both  $u$  and  $v$  are additive this is possible by choosing elements  $x^0, x^1 \in X$  such that  $x^1 \succ x^0$ , and by defining  $u(x^0) = v(x^0) = 0$  and  $u(x^1) = v(x^1) = 1$ . Solving for  $a$  and  $b$  in the solution (14) of Theorem 1 gives the results  $a = 1$  and  $b = 0$ ; therefore

$$u = v. \quad (68)$$

In other words, if  $u$  and  $v$  are standardized a priori to assume values zero and one at identical points in  $X$ , they must be identical everywhere.

Using the same standardizations in the case where  $u$  is multiplicative and  $v$  is additive gives the solutions  $b = 1$  and  $a = (\text{sgn } k) \ln(1+k)$ . Substituting  $a$  and  $b$  in the solution (26) of Theorem 2 gives

$$1+ku = (1+k)^v. \quad (69)$$

Figure 1 gives examples of such functions, which only depend on the "interaction" parameter  $k$  in the multiplicative model. In the literature  $k$  typically is found to lie somewhere between  $-.95$  and  $-.1$  (see Keeney and Raiffa, 1976). For such negative values  $h$  is concave, which is interpreted as risk aversion in value. For  $k \rightarrow -1$  the risk aversion increases, i.e., the function becomes more concave. For  $k \rightarrow 0$  the function approaches  $u = v$ , which corresponds to the result that for  $k \rightarrow 0$   $u$  goes over into the additive model. For  $k > 0$  the function  $h$  becomes convex with increasing convexity (risk proneness) as  $k$  gets larger.

The range between 0 and 1 is of particular interest for analyzing the curvature of  $h$ , since typically both  $u$  and  $v$  would be standardized by selecting  $x^0$  as the worst alternative and  $x^1$  as the best. Figure 2 shows a blowup of Figure 1 in the local range between 0 and 1. For  $k$  values between  $-.5$  and  $+1$  the functional relationship between  $u$  and  $v$  is almost linear, but for  $k = -.99$  concavity (risk aversion) is substantial, and for  $k = 50$  convexity (risk proneness) is significant.

---

Insert Figures 1 and 2 about here

---

In the case where  $u$  is additive and  $v$  is multiplicative the above standardization conventions cannot be used since  $v > 0$  everywhere. However, applying the same conventions to  $u$  and defining  $v(x^0) = 1$  and  $v(x^1) = e$  gives the convenient solutions  $a = 1$  and  $b = 0$ . Using these values in (52), the solution of Theorem 3 gives


$$u = \ln v. \quad (70)$$

When both  $u$  and  $v$  are multiplicative, standardizations are constrained by the fact that  $(1+ku)$  and  $v$  must obtain zero values at identical points. This fixes a point of  $u$  as a function of  $v$  and makes it necessary to standardize separately for the cases where  $k$  is positive and where  $k$  is negative. If  $k$  is positive, the following conventions will be used:  $u(x^0) = 0$ ,  $v(x^0) = 1$ ;  $u(x^1) = 1$ ,  $v(x^1) = e$ . These are the same conventions used above, but there is an additional restriction:  $x^0$  must not be a natural zero point (i.e., a zero multiplier), and for any zero multiplier  $z^0$ ,  $x^1 > x^0 > z^0$ . These conventions give solutions  $b = 1$  and  $a = \ln(1+k)$ . Consequently the solution (67) of Theorem 4 reduces to

$$1+ku = (\operatorname{sgn} v) |v|^{\ln(1+k)}, \quad k > 0. \quad (71)$$

FIGURE 1

GLOBAL PROPERTIES OF FUNCTIONS RELATING  $u$  AND  $v$  IF  $u$  IS  
MULTIPLICATIVE AND  $v$  IS ADDITIVE (Standardization conventions as in text).



**LOCAL PROPERTIES OF FUNCTION RELATING  $u$  AND  $v$  IF  $u$  IS MULTIPLICATIVE AND  $v$  IS ADDITIVE (Standardization conventions as in text)**

$$(1+ku) = (1+k)^v$$




Figure 3 gives examples of this functional relationship which again only depends on  $k$ .  $v = 0$  is in an inflection point for all  $k$ , and the positive and negative sides are mirror images up to a multiplicative transformation. This fact follows from the character of negative multipliers which should maintain the essential properties of the utility function in its negative and positive part.  $k = e - 1$  gives the linear relationship. For larger  $k$ ,  $h$  is concave in negative values of  $v$  and convex in positive ones. For smaller  $k$  this trend reverses. A final observation is the natural limit of the concavity of  $h$  when  $k$  tends towards zero. The dotted line represents this limit, which is in fact the function  $u = \ln v$ , as would be predicted since  $u$  becomes additive if  $k$  goes to zero.

---

Insert Figure 3 about here

---

Figure 4 gives again the blowup for the standard range of  $u$  between 0 and 1. Here it appears that  $k$  values between 1 and 5 tend to produce close to linear functions.

---

Insert Figure 4 about here

---

When  $k$  is negative the following standardization conventions will be used:  $u(x^0) = 0$ ,  $v(x^0) = -1$ ;  $u(x^1) = -1$ ,  $v(x^1) = -e$ . These are similar to the conventions for positive  $k$ , but applied to the negative ranges of  $u$  and  $v$ . Here the additional assumption must be made that  $x^0$

FIGURE 3

GLOBAL PROPERTIES OF FUNCTIONS RELATING  $u$  AND  $v$  IS BOTH ARE  
MULTIPLICATIVE (Standardization conventions for  $k > 0$  as in text)





FIGURE 4

LOCAL PROPERTIES OF FUNCTIONS RELATING  $u$  AND  $v$  IF BOTH ARE  
MULTIPLICATIVE (Standardization conventions for  $k > 0$  as in text)

$$1 + ku = (\text{sgn } v) |v|^{\ln(1+k)}$$

$$k \rightarrow 0, u = \ln v$$



is not a zero multiplier, and that for any zero multiplier  $z^0$   $z^0 > x^0 > x^1$ . Using these standardization conventions, the solutions for  $a$  and  $b$  are  $b = 1$  and  $a = \ln(1-k)$ . Substituting these solutions into (53), the solution to Theorem 5 gives

$$1+ku = -(\text{sgn } v) |v|^{\ln(1-k)}, k < 0 \quad (72)$$

Figures 5 and 6 present examples of this functional relationship both for the global range of  $u$  and  $v$  and the local range within the standardized values.  $v = 0$  is again an inflection point, and the positive and negative segments are mirror images up to a multiplicative transformation. In this case, however,  $h$  is convex (risk seeking) in the negative values of  $k$  and concave (risk prone) in the positive values. The local curvature of  $h$  within the standardized range is close to linear for most of the range of negative  $k$ 's.

---

Insert Figures 5 and 6 about here

---

### Behavioral and Applied Implications

Behavioral implications. To explore the behavioral meaning of the mathematical results it is helpful to consider a simple commodity bundle evaluation example. Let each  $X_i$  denote a commodity with unit price  $p_i$ .  $x = (x_1, x_2, \dots, x_i, \dots, x_n)$  is a commodity bundle with amounts  $x_i$ . The market price for such a bundle would be

$$p(x) = \sum_{i=1}^n p_i x_i. \quad (73)$$

GLOBAL PROPERTIES OF FUNCTIONS RELATING  $u$  AND  $v$  IF BOTH ARE  
MULTIPLICATIVE (Standardization conventions for  $k < 0$  as in text)





FIGURE 6

LOCAL PROPERTIES OF FUNCTIONS RELATING  $u$  AND  $v$  IF BOTH ARE  
MULTIPLICATIVE (Standardization conventions for  $k > 0$  as in text)



(73) is also a simple additive value model in which preferences for (riskless) commodity bundles are dictated by their respective prices. The implications of linear value functions  $v_i = p_i x_i$  and linear indifference curves are obvious.

Assuming that an analyst only knows that (73) is an appropriate conjoint measurement value representation for some decision maker and that the v. Neumann and Morgenstern utility function  $u$  is additive, theorem 1 implies the strong results:

$$u(x) = \sum_{i=1}^n p_i x_i, \quad (74)$$

$$u_i(x_i) = p_i x_i. \quad (75)$$

That, of course, means that the decision maker is risk neutral.

To extend this line of argument to nonlinear value functions, assume that all  $v_i$ 's are marginally decreasing as described by a logarithmic function

$$v_i(x_i) = \ln x_i, \quad (76)$$

and that  $v$  is additive. Assuming again that  $u$  is additive implies

$$u_i = \ln x_i. \quad (77)$$

The single attribute utility functions in (77) appear risk averse, but risk aversion in this particular context means nothing more than marginally

decreasing value. There is absolutely no element of the colloquial notion of "avoiding to take chances" implied by the shape of the  $u_i$ 's. On the contrary, I would consider a decision maker whose preferences follow (76) and (77) risk neutral in value.

This risk neutrality in value comes, in fact, as no surprise considering the structural assumption behind the additivity of  $u$ . This assumption, called the marginality assumption (Fishburn, 1965), requires the decision maker to be indifferent among all gambles with identical single attribute (marginal) probability distributions. To illustrate, consider the following two gambles for commodity bundles:

|             | GAMBLE 1 |       | GAMBLE 2 |       |
|-------------|----------|-------|----------|-------|
|             | HEADS    | TAILS | HEADS    | TAILS |
| GASOLINE    | 16 G     | 0     | 0        | 16 G  |
| GROUND BEEF | 10 P     | 0     | 10 P     | 0     |

According to the marginality assumption these two gambles should be indifferent. But an experimental study (v. Winterfeldt, 1979) revealed that subjects generally prefer the right option, because it has more balanced outcomes. Similar results were obtained by Delbeke and Fauville (1974) and in applied studies (Keeney and Raiffa, 1976). This is, of course, a form of risk aversion but to distinguish it from the usual single attribute risk aversion it has been called multiattribute risk aversion (see also Richard, 1975). The marginality assumption is therefore equivalent to multiattribute risk neutrality, and single attribute risk neutrality in value. It does not, however, require linear utility

functions as example (77) shows.

Disentangling risk attitude from marginal value considerations becomes more interesting if we consider the riskless model  $v(x) = \sum_{i=1}^n \ln x_i$  in connection with a multiplicative utility function. The standardized form of Theorem 2 implies

$$1+ku(x) = (1+k) \sum_{i=1}^n \ln x_i. \quad (78)$$

Setting all  $x_j = x_j^0$  except for some  $i$  gives

$$1+ku_i(x_i) = (1+k) \ln x_i. \quad (79)$$

Now assume further than  $k = e-1$ , which yields

$$u_i = \frac{1}{e-1} x_i - \frac{1}{e-1}. \quad (80)$$

which is a linear utility function. This example shows that it is theoretically possible to have linear utility functions, not because the decision maker is risk neutral, but because his risk proneness compensates his marginal decreasing value function. The example also shows that the marginality tests could conceivably be violated in spite of linear utility functions. Or, in other words, a decision maker may be multivariate risk averse (prone) in spite of single attribute risk neutrality. Of course, this single attribute risk neutrality would only be an appearance when  $u$  is multiplicative and  $v$  is additive, since it may be based on strong risk aversion or proneness in value.

Multiattribute risk aversion and proneness can be related to the interaction parameter  $k$  in the multiplicative model (see Keeney and Raiffa, 1976). If  $k < 0$ , the decision maker must be multiattribute risk averse, i.e., he would prefer the more balanced gamble in the marginality test. If  $k > 0$ , the decision maker must be multiattribute risk prone, i.e., he would prefer the more imbalanced gamble in the marginality test.

Theorems 3 and 4 do not contribute substantively to the behavioral understanding of riskless and risky evaluation phenomena. The case where  $u$  is additive and  $v$  is multiplicative (in the non-genuine sense of having no null zones or sign reversals) appears to be very rare. In fact, from a strictly measurement theoretic point of view it cannot happen at all, since additivity of  $u$  implies the existence of an additive  $v$ . In the conjoint measurement sense a multiplicative  $v$  without null zones or sign reversals is indistinguishable from an additive  $v$ . To maintain that  $v$  is multiplicative in such a case requires a priori reasoning which lies outside the arguments of conjoint measurement theory. For example, in evaluating cars, the attributes safety and performance may be considered multiplicative factors a priori, even if there are not preference reversals or null multipliers in the attribute ranges under consideration. Such multiplicative versions of a theoretically additive conjoint measurement model may be preferable for face validity reasons. Consider the case of a non-genuine multiplicative  $v$  where all  $v_i$ 's are exponential in the form  $v_i(x_i) = e^{x_i}$ , and where  $u$  is additive. By Theorem 3

$$u = \ln x = \sum_{i=1}^n x_i. \quad (81)$$

(81) demonstrates again the possibility of a linear utility function with curvilinear value functions.

Genuinely multiplicative value and utility functions are very rare and if they occur they are striking phenomena as Krantz et al note (Krantz et al, 1971). The behavioral interpretations for functional relationships between  $u$  and  $v$  in this case are less clear than in the previous cases. Assume that  $v$  is multiplicative in the form

$$v(x) = \prod_{i=1}^n x_i. \quad (82)$$

If  $u$  is also multiplicative, the standardized solutions of Theorem 4 implies

$$u_i(x_i) = (\text{sgn } k) \frac{1}{k} (\text{sgn } x_i) |x_i|^{\ln[1 + (\text{sgn } k)k]} \quad (83)$$

which is risk neutral if  $k = e - 1$ , risk prone in positive  $x_i$  if  $k > e-1$  and risk averse in positive  $x_i$  if  $k < e-1$ . The negative  $x_i$ 's have the opposite risk attitude when compared with the positive  $x_i$ 's.

Applied implications. The main applied argument for using Theorems 1-4 is that  $v$  can be constructed with simpler methods than  $u$ .  $u$  can be obtained from  $v$  on the basis of a priori reasoning about the aggregation form of  $u$  and  $v$ , and on the basis of a few risky questions to assess  $k$ .

Why is  $v$  easier to construct than  $u$ ? In (conjoint measurement) theory a special sequence of indifference judgments is required to construct  $v$ . This procedure is called "dual standard sequence" or "lock step procedure" (see Krantz, et al, 1971 and Keeney and Raiffa, 1976). This procedure is not substantially simpler than the lottery methods required to construct  $u$ . But simple rating and weighting techniques, notably Edwards' SMART procedure (1977), can approximate dual standard sequences, since this involves similar cognitive processes. It would be much more difficult to justify a SMART approximation of  $u$ , since risk attitude and cognitive processes particular to lottery methods are involved in the direct assessment of  $u$ .

Assuming that SMART or some similar methods are fair or good approximations of  $v$ , Theorems 1-4 provide a simple basis of transforming  $v$  into  $u$  and of crosschecking the construction of value and utility functions. If the considerations of Theorem 1 apply, SMART can be used directly to take expectations, since  $u = v$ . If the conditions of Theorem 2 are met, the analyst has to assess the interaction parameter  $k$  in addition to  $v$ . Often this can be done by a priori reasoning about multiattribute risk attitude (to determine the sign of  $k$ ) and by a few exploratory questions about the degree of multivariate risk attitude (to determine the size of  $k$ ). Sometimes translating  $v$  into  $u$  may not even be worth the effort, e.g., if  $k$  is between  $-.5$  and  $+1$ . Figures 1 and 2 show that in such cases the exponential transformations are almost linear, and expected utilities would be very difficult to distinguish from expected values taken over  $v$ .

Similar methods can be applied in the cases where the conditions of Theorems 3 and 4 are met. In all cases it would be good practice to construct both  $u$  and  $v$  and cross check their implications, when the transformation of  $v$  is assessed.

This last argument leads up to a development of methods for performing sensitivity analyses, based on Theorems 1-4. In particular, these theorems give some simple deterministic bounds for modelling errors, i.e., from using an additive SMART type model when the true model is multiplicative. By now, the steps to perform such sensitivity analysis should be obvious. Further experimental and numerical studies of the functional forms in the four theorems may prove useful for applications of decision analysis.

## REFERENCES

1. J. AZCEL, Lectures on Functional Equations and Their Applications, Academic Press, New York, 1966.
2. L. DELBEKE and J. FAUVILLE, "An Empirical Test of Fishburn's Additivity Axiom," Acta Psychologica 38, 1-20 (1974).
3. J.S. DYER and R.K. SARIN, "Measurable Multiattribute Value Functions," Discussion Paper No. 66, Management Science Study Center, University of California, Los Angeles, 1977.
4. W. EDWARDS, "How to Use Multiattribute Utility Measurement for Social Decision Making," IEEE Transactions on Systems, Man, and Cybernetics 7, 326-340 (1977).
5. P.C. FISHBURN, Utility Theory for Decision Making, Wiley, New York, 1970.
6. P.C. FISHBURN and R.L. KEENEY, "Generalized Utility Independence and Some Implications", Operations Research 23, 928-940 (1974).
7. R.L. KEENEY and H. RAIFFA, Decisions with Multiple Objectives: Preferences and Value Tradeoffs, Wiley, New York, 1976.
8. D.H. KRANTZ and A. TVERSKY, "Conjoint-Measurement Analysis of Composition Rules in Psychology," Psychological Review 78, 151-169 (1970).
9. D.H., KRANTZ, R.D., LUCE, P. SUPPES, and A. TVERSKY, Foundations of Measurement, Vol. I, Academic Press, New York, 1971.
10. J. VON NEUMANN and O. MORGENSTERN, Theory of Games and Economic Behavior, 2nd ed., Princeton University Press, Princeton, N.J., 1947.
11. R.A. POLLAK, "Additive von Neumann and Morgenstern Utility Functions." Econometrica 35, 485-494 (1967).
12. H. RAIFFA, "Preferences for Multi-Attributed Alternatives," Memorandum No. 5868, The Rand Corporation, Santa Monica, 1969.
13. S.F. RICHARD, "Multivariate Risk Aversion, Utility Independence, and Separable Utility Functions," Management Science 22, 12-21 (1975).
14. L.J. SAVAGE, The Foundations of Statistics, Wiley, New York, 1954.
15. D. v. WINTERFELDT, "Additivity and Expected Utility in Risky Multi-attribute Preferences," submitted to Journal of Mathematical Psychology, 1979.

| REPORT DOCUMENTATION PAGE  |  | READ INSTRUCTIONS<br>BEFORE COMPLETING FORM                 |
|--|--|---|
| 1. REPORT NUMBER<br>001595   | 2. GOVT ACCESSION NO.  | 3. RECIPIENT'S CATALOG NUMBER                               |
| 4. TITLE (and Subtitle)<br><br>Functional Relationships Between Risky and Riskless Multiattribute Utility Functions  | 5. TYPE OF REPORT & PERIOD COVERED<br>Technical<br>12/14/78 - 12/15/79 |   |
|  | 6. PERFORMING ORG. REPORT NUMBER<br>79 - 3                             |   |
| 7. AUTHOR(s)<br>Detlof von Winterfeldt   | 8. CONTRACT OR GRANT NUMBER(s)<br>N00014-79-C-0038                     |   |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS<br>Social Science Research Institute<br>University of Southern California<br>Los Angeles, CA 90007   |  | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS |
| 11. CONTROLLING OFFICE NAME AND ADDRESS<br>Office of Naval Research - Dept. of Navy<br>800 N. Quincy Street<br>Arlington, VA 22217   | 12. REPORT DATE<br>December, 1979                                      |   |
|  | 13. NUMBER OF PAGES<br>34  |   |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)<br>Office of Naval Research<br>Research Branch Office<br>1030 East Green Street<br>Pasadena, CA 91101  | 15. SECURITY CLASS. (of this report)<br><br>unclassified               |   |
|  | 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE                             |   |
| 16. DISTRIBUTION STATEMENT (of this Report)<br><br>Approval for public release, distribution unlimited   |  |   |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)   |  |   |
| 18. SUPPLEMENTARY NOTES  |  |   |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number)<br>conjoint measurement theory                      additive & multiplicative utility functions<br>expected utility theory                              value functions<br>functional equations<br>risky and riskless utility functions   |  |   |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number)<br>Expected utility theory and conjoint measurement theory form two major classes of models and assessment procedures to construct multiattribute utility functions. In conjoint measurement theory a value function $v$ is constructed which preserves preferences among riskless multiattributed outcomes. The risky utility function $u$ , constructed in the framework of expected utility theory, also preserves such riskless preferences. In addition, $u$ is an appropriate guide for decisions under uncertainty since its expectation preserves risky preferences among gambles. |  |   |

unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

Since both  $u$  and  $v$  are order preserving functions, they must be related by a strictly increasing transformation. However,  $u$  and  $v$  need not coincide or be related through any special functional forms, unless some simple decomposition forms are assumed. More restricted functional relationships obtain, if  $u$  and  $v$  are assumed to be either additive or multiplicative. In particular,  $u$  can be shown to be linearly, logarithmically, or exponentially related to  $v$ , depending on which function is additive and which is multiplicative. The paper proves such functional relationships based on the theory of functional equations, and techniques are described to assess the parameters of these functions. The results are discussed from a behavioral standpoint of interpreting the form and shape of multiattribute utility functions and from a practical standpoint of simplifying multiattribute utility assessment.

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

CONTRACT DISTRIBUTION LIST  
(Unclassified Technical Reports)

|   |           |
|---|-----------|
| Director<br>Advanced Research Projects Agency<br>Attention: Program Management Office<br>1400 Wilson Boulevard<br>Arlington, Virginia 22209 | 2 copies  |
| Office of Naval Research<br>Attention: Code 455<br>800 North Quincy Street<br>Arlington, Virginia 22217                                     | 3 copies  |
| Defense Documentation Center<br>Attention: DDC-TC<br>Cameron Station<br>Alexandria, Virginia 22314  | 12 copies |
| DCASMA Baltimore Office<br>Attention: Mr. K. Gerasim<br>300 East Joppa Road<br>Towson, Maryland 21204                                       | 1 copy    |
| Director<br>Naval Research Laboratory<br>Attention: Code 2627<br>Washington, D.C. 20375   | 6 copies  |

SUPPLEMENTAL DISTRIBUTION LIST  
(Unclassified Technical Reports)

Department of Defense

Director of Net Assessment  
Office of the Secretary of Defense  
Attention: MAJ Robert G. Gough, USAF  
The Pentagon, Room 3A930  
Washington, DC 20301

Assistant Director (Net Technical Assessment)  
Office of the Deputy Director of Defense  
Research and Engineering (Test and  
Evaluation)  
The Pentagon, Room 3C125  
Washington, DC 20301

Director, Defense Advanced Research  
Projects Agency  
1400 Wilson Boulevard  
Arlington, VA 22209

Director, Cybernetics Technology Office  
Defense Advanced Research Projects Agency  
1400 Wilson Boulevard  
Arlington, VA 22209

Director, ARPA Regional Office (Europe)  
Headquarters, U.S. European Command  
APO New York 09128

Director, ARPA Regional Office (Pacific)  
Staff CINCPAC, Box 13  
Camp H. M. Smith, Hawaii 96861

Dr. Don Hirta  
Defense Systems Management School  
Building 202  
Ft. Belvoir, VA 22060

Chairman, Department of Curriculum  
Development  
National War College  
Ft. McNair, 4th and P Streets, SW  
Washington, DC 20319

Defense Intelligence School  
Attention: Professor Douglas E. Hunter  
Washington, DC 20374

Vice Director for Production  
Management Office (Special Actions)  
Defense Intelligence Agency  
Room 1E963, The Pentagon  
Washington, DC 20301

Command and Control Technical Center  
Defense Communications Agency  
Attention: Mr. John D. Hwang  
Washington, DC 20301

Department of the Navy

Office of the Chief of Naval Operations  
(OP-951)  
Washington, DC 20450

Office of Naval Research  
Assistant Chief for Technology (Code 200)  
800 N. Quincey Street  
Arlington, VA 22217

Office of Naval Research (Code 130)  
800 North Quincey Street  
Arlington, VA 22217

Office of Naval Research  
Naval Analysis Programs (Code -91)  
800 North Quincey Street  
Arlington, VA 22217

Office of Naval Research  
Operations Research Programs (Code 434)  
800 North Quincy Street  
Arlington, VA 22217

Office of Naval Research  
Information Systems Program (Code 437)  
800 North Quincy Street  
Arlington, VA 22217

Director, ONR Branch Office  
Attention: Dr. Charles Davis  
536 South Clark Street  
Chicago, IL 60605

Director, ONR Branch Office  
Attention: Dr. J. Lester  
495 Summer Street  
Boston, MA 02210

Director, ONR Branch Office  
Attention: Dr. E. Gloye  
1030 East Green Street  
Pasadena, CA 91106

Director, ONR Branch Office  
Attention: Mr. R. Lawson  
1030 East Green Street  
Pasadena, CA 91106

Office of Naval Research  
Scientific Liaison Group  
Attention: Dr. M. Bertin  
American Embassy - Room A-407  
APO San Francisco 96503

Dr. A. L. Slafkosky  
Scientific Advisor  
Commandant of the Marine Corps (Code RD-1)  
Washington, DC 20380

Headquarters, Naval Material Command  
(Code 0331)  
Attention: Dr. Heber G. Moore  
Washington, DC 20360

Dean of Research Administration  
Naval Postgraduate School  
Attention: Patrick C. Parker  
Monterey, CA 93940

Superintendent  
Naval Postgraduate School  
Attention: R. J. Roland, (Code 5281)  
C3 Curriculum  
Monterey, CA 93940

Naval Personnel Research and Development  
Center (Code 305)  
Attention: LCDR C'Bar  
San Diego, CA 92152

Navy Personnel Research and Development  
Center  
Manned Systems Design (Code 311)  
Attention: Dr. Fred Muckler  
San Diego, CA 92152

Naval Training Equipment Center  
Human Factors Department (Code N215)  
Orlando, FL 32813

Naval Training Equipment Center  
Training Analysis and Evaluation Group  
(Code N-207)  
Attention: Dr. Alfred F. Snodde  
Orlando, FL 32813

Director, Center for Advanced Research  
Naval War College  
Attention: Professor C. Lewis  
Newport, RI 02840

Naval Research Laboratory  
Communications Sciences Division (Code 541)  
Attention: Dr. John Shore  
Washington, DC 20375

Dean of the Academic Departments  
U.S. Naval Academy  
Annapolis, MD 21402

Chief, Intelligence Division  
Marine Corps Development Center  
Quantico, VA 22134

#### Department of the Army

Deputy Under Secretary of the Army  
(Operations Research)  
The Pentagon, Room 2E611  
Washington, DC 20315

Director, Army Library  
Army Studies (ASDIRS)  
The Pentagon, Room 1A534  
Washington, DC 20310

U.S. Army Research Institute  
Organizations and Systems Research Laboratory  
Attention: Dr. Edgar M. Johnson  
5001 Eisenhower Avenue  
Alexandria, VA 22333

Director, Organizations and Systems  
Research Laboratory  
U.S. Army Institute for the Behavioral  
and Social Sciences  
5001 Eisenhower Avenue  
Alexandria, VA 22333

Technical Director, U.S. Army Concepts  
Analysis Agency  
8100 Woodmont Avenue  
Bethesda, MD 20814

Director, Strategic Studies Institute  
U.S. Army Combat Developments Command  
Carlisle Barracks, PA 17013

Commandant, Army Logistics Management Center  
Attention: DRXMC-LS-SCAD (ORSA)  
Ft. Lee, VA 23801

Department of Engineering  
United States Military Academy  
Attention: COL Ar F. Grum  
West Point, NY 10996

Marine Corps Representative  
U.S. Army War College  
Carlisle Barracks, PA 17013

Chief, Studies and Analysis Office  
Headquarters, Army Training and Doctrine  
Command  
Ft. Monroe, VA 23351

Commander, U.S. Army Research Office  
(Durham)  
Box CM, Duke Station  
Durham, NC 27706

#### Department of the Air Force

Assistant for Requirements Development  
and Acquisition Programs  
Office of the Deputy Chief of Staff for  
Research and Development  
The Pentagon, Room 4C331  
Washington, DC 20330

Air Force Office of Scientific Research  
Life Sciences Directorate  
Building 410, Bolling AFB  
Washington, DC 20332

Commandant, Air University  
Maxwell AFB, AL 36112

Chief, Systems Effectiveness Branch  
Human Engineering Division  
Attention: Dr. Donald A. Topmiller  
Wright-Patterson AFB, OH 45433

Deputy Chief of Staff, Plans, and  
Operations  
Directorate of Concepts (AR/MOCCC)  
Attention: Major R. Linhard  
The Pentagon, Room 4D 11-7  
Washington, DC 20330

Director, Advanced Systems Division  
(AFHRL/AS)  
Attention: Dr. Gordon Eckstrand  
Wright-Patterson AFB, OH 45433

Commander, Rome Air Development Center  
Attention: Mr. John Atkinson  
Griffis AFB  
Rome, NY 13440

IRD, Rome Air Development Center  
Attention: Mr. Frederic A. Dion  
Griffis AFB  
Rome, NY 13440

HQS Tactical Air Command  
Attention: LTCOL David Bianich  
Langley AFB, VA 23665

#### Other Government Agencies

Chief, Strategic Evaluation Center  
Central Intelligence Agency  
Headquarters, Room 2G24  
Washington, DC 20505

Director, Center for the Study of  
Intelligence  
Central Intelligence Agency  
Attention: Mr. Dean Moor  
Washington, DC 20505

Mr. Richard Heuer  
Methods & Forecasting Division  
Office of Regional and Political Analysis  
Central Intelligence Agency  
Washington, DC 20505

Office of Life Sciences  
Headquarters, National Aeronautics and  
Space Administration  
Attention: Dr. Stanley Deutsch  
600 Independence Avenue  
Washington, DC 20546

#### Other Institutions

Department of Psychology  
The Johns Hopkins University  
Attention: Dr. Alphonse Chapanis  
Charles and 34th Streets  
Baltimore, MD 21218

Institute for Defense Analyses  
Attention: Dr. Jesse Orlansky  
400 Army Navy Drive  
Arlington, VA 22202

Director, Social Science Research Institute  
University of Southern California  
Attention: Dr. Ward Edwards  
Los Angeles, CA 90007

Perceptronics, Incorporated  
Attention: Dr. Amos Freedly  
6071 Varial Avenue  
Woodland Hills, CA 91364

Stanford University  
Attention: Dr. R. A. Howard  
Stanford, CA 94305

Director, Applied Psychology Unit  
Medical Research Council  
Attention: Dr. A. D. Baddeley  
15 Chaucer Road  
Cambridge, CB 2EF  
England

Department of Psychology  
Brunel University  
Attention: Dr. Lawrence D. Phillips  
Uxbridge, Middlesex UB8 3PH  
England

Decision Analysis Group  
Stanford Research Institute  
Attention: Dr. Miley W. Markhofer  
Menlo Park, CA 94025

Decision Research  
1201 Oak Street  
Eugene, OR 97401

Department of Psychology  
University of Washington  
Attention: Dr. Lee Roy Beach  
Seattle, WA 98195

Department of Electrical and Computer  
Engineering  
University of Michigan  
Attention: Professor Kan Chen  
Ann Arbor, MI 94135

Department of Government and Politics  
University of Maryland  
Attention: Dr. Davis B. Bobrow  
College Park, MD 20747

Department of Psychology  
Hebrew University  
Attention: Dr. Amos Tversky  
Jerusalem, Israel

Dr. Andrew P. Sage  
School of Engineering and Applied  
Science  
University of Virginia  
Charlottesville, VA 22901

Professor Raymond Tanter  
Political Science Department  
The University of Michigan  
Ann Arbor, MI 48109

Professor Howard Raiffa  
Morgan 302  
Harvard Business School  
Harvard University  
Cambridge, MA 02163

Department of Psychology  
University of Oklahoma  
Attention: Dr. Charles Gettys  
455 West Lindsey  
Dale Hall Tower  
Norman, OK 73069

Institute of Behavioral Science #3  
University of Colorado  
Attention: Dr. Kenneth Hammond  
Room 201  
Boulder, Colorado 80309

Decisions and Designs, Incorporated  
Suite 600, 8400 Westpark Drive  
P.O. Box 907  
McLean, Virginia 22101

END

9-87

Dtic